arXiv:astro-ph/0003348v1 23 Mar 2000

Key aspects in the photometric monitoring of δ Sct stars

Ennio Poretti
Osservatorio Astronomico di Brera, Via E. Bianchi 46, I-23807 Merate, Italy

ABSTRACT

The MONS experiment will be able to verify some of the most important topics in the asteroseismologic studies of δ Sct stars. This research will benefit not only from the large increase in the number of detected frequencies, but also from the possibility to perform mode identification by means of very precise two–colour photometry. To do that, an appropriate frequency resolution, comparable to or better than what is currently achieved by ground–based observations, is necessary and hence two runs are recommended. To avoid over-scheduling of the Main Camera, a possible combination of observations from the Main Camera and from the Star Trackers is proposed. As a result, two key-objects as FG Vir and BH Psc could be adequately monitored, affording the possibility to enter into details of amplitude variations. As a by-product, the same results could be obtained on two γ Dor stars.

1. Introduction

In the last two decades a great observational effort has been undertaken to solve the light curves of δ Sct stars. These complicated variables, located on or just above the Zero–Age–Main–Sequence, constitute the lower limit of the classical instability strip. The presence of convective zones makes these stars very interesting from the evolutionary point of view, since various stages, from pre–main sequences to giant, can be observed.

Photometric monitoring is the most practiced approach to study their properties and several stars have been deeply investigated; for a review, see Poretti (2000). However, the spectroscopic approach tells us that many modes that are not photometrically detectable are actually excited (Mantegazza 2000).

The observed frequencies are between 60 and 405 μHz and multiperiodicity is very common; rotation acts as an important factor in the increase of the number of excited modes. Only a few stars show a monoperiodic behaviour above the current limit of the detectable amplitude from ground, i.e. $\sim$1 mmag. Of course, the number of detected modes is strongly dependent on the accuracy, the number and the sampling of the measurements.

The observed modes are in the domain of pressure (p) modes; there is no observational evidence that gravity (g) modes are excited in δ Sct stars, even if some cases suggest they might be. At the moment, g-modes seem to be present only in the γ Dor stars, which in turn do not show p–modes.

How the photometric results can be used to identify modes (i.e. to classify the oscillation in terms of quantum numbers n, ℓ and m) is discussed by Garrido (2000).

2. The state–of–art in the study of δ Sct stars

Several approaches have been used in the past to attack the problem of the detection of excited modes in their light curves. The debate on the nature of the variations, multiperiodic or irregular, occupied the years from 1974 to 1980. At that time the observations consisted in time–series collected over a few nights, a practice that had been borrowed from the study of high amplitude δ Sct stars. In the years that followed, a number of teams observed more carefully δ Sct stars and clearly evidenced their multiperiodicity, detecting stable frequencies and obtaining very satisfactory fits. It appeared also evident that nonradial modes are excited. The number of frequencies is so high that the aliases at 1 cd^{-1} become a serious problem in the mode detection. For that, multisite campaigns were planned and the best results now available have been obtained from these ground–based observations.

The effort made by the observers these last years allowed us to detect a large number of frequencies in many stars: we will recall here the cases of FG Vir, XX Pyx, 4 CVn and BH Psc. By means of multisite campaigns we are able to detect terms with amplitude less than 1 mmag. The mode identification techniques are based on both the phase shifts and amplitude ratios of the light and colour curves and the synergic approach performed by considering spectroscopic curves. Their full exploitation should guarantee an important role for our researches in stellar physics.

Figure 1 shows the frequency content of three well-studied pulsators in which all the terms with a half-amplitude larger than 1 mmag are detected. It appears that their content is completely different: high frequencies only for XX Pyx, low frequencies only for 4 CVn, two groups of intermediate frequency values for FG Vir. The combination terms are also reported in the upper part of each panel; 4 CVn displays both $f_i - f_j$ and $f_i + f_j$ terms, located right and left of the bunch of independent frequencies. FG Vir displays only three combination modes in the same region of the independent frequencies. XX Pyx displays only a $2f$ harmonic, at 76.2 cd^{-1}.

The obvious conclusion is that δ Sct stars are very complicated pulsators and that a general recipe cannot be applied to predict the range of the excited modes or the importance of combination terms. However, a curious similarity in the frequency content was noted by Mantegazza & Poretti (1999) comparing 4 CVn to HD 2724: at least 7 frequencies have almost the same value.

3. The amplitude variations

The variations of the amplitudes have been detected when a δ Sct star has been re–observed in order to investigate some unclear facts: a comparison of the mode amplitudes in the two



Fig. 1.— In δ Sct stars the different distributions of the independent frequencies (lower line in each panel) and of the combination terms (upper line in each panel; XX Pyx shows only a harmonic term at 76.2 cd^{-1}, not displayed here) are evident when comparing the cases of XX Pyx, 4 CVn and FG Vir. Amplitudes are in mmag unit.

different seasons often evidences strong differences. Even if the data sampling can affect the reliability of these conclusions, some results seem to be very much in favour of a physical variability (Breger 2000). These variations can be seen as a further complication, even if we can argue that the disappearance or the damping of some terms can enhance or make discernible other modes, increasing the number of known frequencies.

Amplitude variations are observed in a large variety of stars, both multiperiodic (XX Pyx, Handler et al. 1997; 4 CVn, Breger 2000; 44 Tau, Poretti et al. 1992) and monoperiodic (28 And, Rodríguez et al. 1998; BF Phe, Poretti et al. 1996). The observations of mode growth in the light curve of a relatively simple pulsator as V663 Cas can clarify what happens in much more complicated cases: some modes can be damped and then re–excited (Poretti et al. 1996). Some observational facts favour this explanation even if the model of a beating between two close frequencies with similar, constant amplitude cannot be ruled out.

BH Psc is another example. It was observed by our group in 1991 (on two occasions), 1994 and 1995 at the European Southern Observatory, Chile. The analysis of the first two campaigns showed a very complex light variability resulting from the superimposition of more than 10 pulsation modes with frequencies between 5 and 12 cd^{-1} and semi–amplitudes between 17 and 3 mmag (Mantegazza, Poretti, & Zerbi 1995). The fit left a high r.m.s. residual 2.3 times greater than the one measured between the two comparison stars. A second photometric campaign was carried out in October and November 1994; we hoped to reveal more terms and to check the stability of their amplitudes. The analysis of the new data allowed us to single out 13 frequencies (Mantegazza, Poretti, & Bossi 1996). They are concentrated in the region 5–11.5 cd^{-1}: this distribution is slightly larger than the one observed for 4 CVn, but they can be considered very similar. However, more terms should be present, with an amplitude below 3 mmag, since about 15% of the variance could not be explained with the detected terms and the noise. Moreover, the standard deviation around the mean value was considerably higher in 1991 than in 1994 (26 mmag against 18 mmag); considerations about the amplitudes of the modes confirmed that the pulsation energy was lower in 1994 than in 1991. Moreover, comparing the 1991 September and 1991 October data it is quite evident that we can detect such variations on a baseline of 1 month or less (see Tab. 1, where the results of a preliminary analysis are shown).

4. Scientific topics to be matched by MONS

The precision in the measurements achievable by MONS will of course allow us to detect terms with amplitudes much smaller than 1 mmag. This in turn will generate a much more detailed knowledge of the frequency content of δ Sct variables. We could then obviously match one of the requests of the asteroseismology theory, i.e. increasing the number of detected terms. This is quite obviuos. However, the values of the amplitudes of these modes are also an important parameter that can help in the theoretical models. The observational evidence is that these amplitudes are

Table 1: Preliminary values for the amplitudes of different terms observed in the light curve of BH Psc

Frequency	Sept. 1991	Oct. 1991	Oct. 1994	Oct. 1995
[cd^{-1}]		[mmag]		
6.214	17.5	17.1	15.0	18.4
9.538	13.8	9.8	1.1	8.2
10.241	9.1	8.2	9.9	15.4
10.561	5.7	16.5	2.3	14.1
8.486	13.3	14.1	0.3	3.1
7.118	10.6	4.6	6.2	8.2
8.788	7.0	4.9	4.4	9.0
11.423	9.1	4.9	6.6	5.7

variable. Can MONS then give us an important clue as regard this aspect ?

To reach the goal, observing twice the same field can be very useful. Doing so, we could:

1. gain in frequency resolution since, as shown in Fig. 1, there are many close terms. It should also be noted that multisite campaigns span more than 50 d and that a single 30-50 d MONS run will not have a sufficient frequency resolution to solve already known close peaks. Owing to the expected large number of peaks, interference between different terms can seriously affect the quality of the results: an unappropriate frequency resolution could be considered a weakness of the MONS results;

2. detect amplitude variations in a very clear way. The absence of alias terms combined with the appropriate frequency resolution will allow a clarification on the nature of the amplitude variations, i.e. beating or intrinsic variations.

5. A plausible solution

In the current configuration, can MONS satisfy these two requirements ?

Among the δ Sct stars, FG Vir is by far the most studied: not only an excellent observational basis is available, but also more and more thorough theoretical models are proposed on the basis of the observed frequencies (Breger et al. 1999). FG Vir (α $12^{\rm h}14^{\rm m}15^{\rm s}$, $\delta - 5^\circ43'00''$, 2000.0) is located very close to β Vir (α $11^{\rm h}50^{\rm m}42^{\rm s}$, $\delta + 1^\circ45'53''$, 2000.0), a F9 V star candidate for searching solar–like oscillations. Then, FG Vir (V=6.1) can be measured not only as a primary target with the Main Camera, but also with the Forward–pointing star tracker (BST or SI-2) when β Vir is

Table 2: A possible observing MONS schedule matching the scientific requirements of the studies on δ Sct stars. See text for explanations.

Main Camera	Frontward ST (BST)	Backward ST (FST)
FG Vir	β Vir II Vir	BH Psc HD 224639 HD 224945
β Vir	FG Vir II Vir	BH Psc HD 224639 HD 224945

measured in the Main Camera. From this lucky coincidence, we can obtain two runs on a well–studied δ Sct star, greatly increasing the number of detected frequencies and saving frequency resolution. Moreover, in both cases, it will be possible to measure II Vir, another δ Sct star close to FG Vir.

Is that the only result ? Certainly not. There would be one more bonus. Looking through the backward–pointing star tracker (FST or SI-1) we could see, in both cases, an interesting field, located at 176° from FG and β Vir. Three intriguing objects are located in this field: namely, BH Psc (α $23^h59^m31^s$, $\delta - 2^\circ50'37''$, 2000.0), i.e. a δ Sct showing strong amplitude variations and HD 224639 and HD 224945, two γ Dor variables. The requirements about frequency resolution and amplitude variations also apply to γ Dor variables.

As a conclusion, it looks possible for MONS to perform an observational study of δ Sct stars in a very powerful way (increase in the number of detected terms, two–colour photometry, sufficiently long baseline allowing frequency resolution and amplitude monitoring) just considering some scheduling precautions and putting an adequate target in the Main Camera.

REFERENCES

Breger, M., 2000, MNRAS, in press

Breger, M., Pamyatnykh, A.A., Pikall, H., Garrido, R., 1999, A&A, 341, 151

Garrido, R., 2000, in "Delta Scuti and Related Stars", Eds. M. Breger and M. Montgomery, ASP Conf. Series, in press

Handler, G., Pikall, H., O'Donoghue, D., Buckley, D.A.H., Vauclair, G., et al., 1997, MNRAS, 286, 303

Mantegazza, L., 2000, in "Delta Scuti and Related Stars", Eds. M. Breger and M. Montgomery, ASP Conf. Series, in press

Mantegazza, L., Poretti, E., 1999, A&A, 348, 139

Poretti, E., 2000, in "Delta Scuti and Related Stars", Eds. M. Breger and M. Montgomery, ASP Conf. Series, in press

Poretti, E., Mantegazza, L., Bossi, M., 1996, A&A, 312, 912

Poretti, E., Mantegazza L., Riboni, E., 1992, A&A, 256, 113

Rodríguez, E., Rolland, A., López-Gonzáles, M.J., Costa, V., 1998, A&A, 338, 905

arXiv:astro-ph/0003348v1 23 Mar 2000

Understanding the γ Doradus Phenomenon using MONS

F.M. Zerbi[1], C. Aerts[2], G. Handler[3], A.B. Kaye[4] and E. Poretti[1]
[1] *Osservatorio Astronomico di Brera, Via Bianchi 46, 23807 Merate (Lc), Italy*
[2] *Instituut voor Sterrenkunde, Katholieke Universiteit Leuven,*
Celestijnenlaan 200 B, B-3001 Leuven, Belgium
[3] *SAAO, PO Box 9, Observatory 7935, South Africa*
[4] *Los Alamos National Laboratory, X-TA, MS B-220, Los Alamos NM 87545, USA*

ABSTRACT

We discuss the results expected from observing γ Dor stars with the MONS satellite. We first describe how MONS Star Trackers will consistently improve the present knowledge of this new class of variables. We then discuss how the γ Dor can be considered possible "bridges" between opacity driven pulsations and stochastically excited solar like oscillations. For this reason the possible inclusion of a γ Dor representative in the MONS main target list is suggested and discussed. Three possible candidates are presented: γ Dor, 9 Aur and BS 2740.

1. A Brief Introduction to γ Doradus Stars

The γ Dor stars have been recently defined (Kaye et al. 1999) as a class of variable stars in the lower-red part of the Cepheid instability strip, undergoing g-mode pulsation. Stars in this group show (multi)periodic photometric and spectroscopic variability with typical time scales between 0.3 and 3 d and typical amplitudes below $0^m.1$. The definition of the class was based on 12 objects contained in a so-called *master list*. The principal physical properties of these stars are reported in Table 1. Several objects suspected to belong to the class have been recently added by Aerts, Eyer & Kestens (1998) and Handler (1999a) upon examination of HIPPARCOS light curves. At the present date the data-base of known γ Dor stars and related objects (Handler, private communication) contains 141 objects divided into *Bona Fide* (12), *Prime* (46) and *Other* (83) candidates. Bona fide objects in the master list are stars for which extensive photometric and spectroscopic data prove that pulsations must be the cause for the variability. Prime and Other γ Dor candidates are stars for which either a moderate or a limited amount of photometric and/or spectroscopic data is available, allowing to suspect or speculate that nonradial pulsation (NRP) is the cause of the observed variations.

The most relevant problems in the observation of γ Dor stars concern the amplitudes and time scales of their variability. Amplitudes are small and the clustering of the typical frequencies around $1\,\mathrm{d}^{-1}$ creates conflicts with the $1\,\mathrm{d}^{-1}$ alias typical of ground-based single-site observations and requires to organize multi-longitude campaigns. At the same time even in these campaigns problems

Star	$b-y$ (mag)	$<V>$ (mag)	ST	$v \sin i$ (kms^{-1})	π (mas)	$[Me/H]$	M_V (mag)	L $[L_\odot]$	$T_{\rm eff}$ (K)	R $[R_\odot]$	M $[M_\odot]$
HD 224945	0.192	6.93	F0 V	55	16.92	-0.30	3.07	5.1	7250	1.43	1.51
γ Dor	0.201	4.25	F0 V	62	49.26	-0.02	2.72	7.0	7200	1.70	1.57
9 Aur	0.217	5.00	F0 V	18	38.14	-0.19	2.89	6.0	7100	1.62	1.52
BS 2740	0.219	4.49	F0 V	40	47.22	-0.15	2.86	6.2	7100	1.64	1.53
HD 62454	0.214	7.15	F1 V	53	11.18	0.16	2.39	9.5	7125	2.02	1.66
HD 68192	0.227	7.16	F2 V	85	10.67	0.05	2.30	10.5	7000	2.20	1.71
HD 108100	0.234	7.14	F2 V	68	12.10	-0.03	2.53	8.5	6950	2.01	1.62
HD 164615	0.226	7.06	F2 IV	66	14.36	0.20	2.82	6.5	7000	1.73	1.53
BS 6767	0.183	6.40	F0 Vn	135	17.44	-0.10	2.59	7.9	7300	1.76	1.61
BS 8330	0.225	6.20	F2 IV	38	19.90	-0.01	2.67	7.4	7000	1.85	1.57
BS 8799	0.181	5.99	kA5	45	25.04	-0.36	2.96	5.7	7375	1.46	1.54
HD 224638	0.198	7.49	F1 Vs	24	12.56	-0.15	2.98	5.5	7200	1.51	1.52

Table 1: Observational parameters of the confirmed γ Dor variables and calculated or inferred basic properties. Table taken from Kaye et al. (1999).

of alignment, improper correction for atmospheric extinction and fluctuation in the instrumental setup often introduce spurious low-frequency terms that have the chance to be misinterpreted as signal. For this reason γ Dor stars will benefit as many other classes of variables of continuous out-of-atmospheric coverage as the one provided by a space mission like MONS.

The interested community has largely discussed if γ Dor stars have any relevance for asteroseismology. These stars pulsate in *g*-modes that in some sense carry more information on the stellar interior than *p*-modes, but the low number of modes excited and the intrinsic uncertainties in the application of standard mode-identification techniques seem to make a seismic interpretation rather difficult. The γ Dor stars give us however two additional possibilities for seismological investigations which other classes of pulsating star cannot provide.

Firstly, the domain of γ Dor stars in the HR diagram partly overlaps with that of the δ Scuti stars (Handler 1999a). It can therefore be suspected that stars showing both types of pulsation could be present. This will allow to constrain the physical properties of a star tightly from the δ Scuti-type *p* modes, which helps in identifying the γ Dor-type *g* modes which can then be used to probe the star's deep interior. Indeed, short-period light variations have already been detected for two γ Dor stars (see Handler 1999b for the first discovery).

Secondly, Guzik et al. (2000) pointed out the major role played by convection in γ Dor pulsation theory. These authors calculated nonadiabatic pulsation properties of evolutionary models lying near the γ Dor instability region in the H-R diagram. The models considered have relatively deep envelope convection zones. By using the Pesnell (1990) non-adiabatic pulsation code these authors calculated the $l = 0, 1, 2$ pulsation frequencies and found unstable high order *g*-modes with frequencies between 4 and 25 μHz (periods $\sim$ 0.4 to 3 days). Furthermore the mode kinetic energy was found to reach a minimum at a frequency of about 11 μHz (period of $\sim$ 1 d), i.e. near the most commonly observed γ Dor period. The frequency spacing between modes with same l and subsequent n was found to be ~ 0.1 d^{-1}, exactly matching the typical frequency spacing observed in multiperiodic γ Dor stars. These results strongly suggest that asteroseismology of γ Dor stars might be easier as it seemed until very recently.

Another interesting outcome of the model calculations originates from the "frozen-in convection" approximation used, in which fluctuations in the convective luminosity are set to zero during the pulsation cycle. Because convection does not adapt to transport the luminosity in this approximation, the radiation is periodically blocked by the high opacity at the convection zone base, resulting in pulsational driving. Therefore the overstable g-modes in γ Dor stars are first maintained by an opacity bump at the basis of the convective layer which has nothing to do with the classical κ/γ mechanism and "tunnel" to the surface through a convective envelope which is "lazy" enough not to adjust and damp them. Second the turbulent motion in the convective envelope could trigger solar-like oscillations as in any other stars of this mass, temperature and luminosity. Indeed the temperature and luminosity range occupied by γ Dor stars completely overlaps with the one in which solar-like oscillations are predicted. In this sense γ Dor stars might represent a sort of "bridge" between overstable pulsation and stochastically generated solar-like oscillations. The MONS primary telescope represents an unique opportunity to investigate such a possible link. For this reason we encourage the MONS science team to consider the inclusion of an object of this kind in the primary target list.

2. Observing γ Doradus stars with the star trackers

MONS star trackers will observe their targets continuously for about 30 days; this will greatly reduce the size of alias peaks. A computed MONS spectral window predicts the strongest alias at .02 mHz ($\sim 1.7\,\mathrm{d}^{-1}$) with an amplitude of 0.2×10^{-6}. This number has to be compared to the typical size of the $1\,\mathrm{d}^{-1}$ alias of ground-based multisite campaigns for γ Dor stars, which is of about 3.1×10^{-1} in amplitude or higher. Most of the available literature on γ Dor stars contains discussions of accurate time-series analysis fighting against aliasing. In this sense the superb quality of MONS spectral window will likely allow major improvements in frequency recognition.

It has often been noticed in published frequency analyses (even among γ Dor candidates) that after subtraction of known signals from the data suprisingly high residuals remained. In the best cases the mean internal error achievable from the ground is 2000 ppm, on the average 4000 ppm is probably a more reliable value; this translates into noise levels around 1000 ppm in the amplitude spectra in the frequency region of interest. At the typical magnitudes of the proposed γ Dor targets MONS Star Trackers will allow clear 3σ detections of signals with amplitudes of the order of a few tens of ppm, providing information more accurate by orders of magnitude with respect to the best observations from Earth. In the case of *prime* or *other* candidates the suspected presence of g-mode pulsation is mostly deduced from survey quality light curves (see for example the left box of Figure 1). The residual scatter in the light curves is typically a few percent ($\sim 10^4$ ppm). MONS STs will improve the measurement accuracy by a factor of at least 100 allowing to confirm the detection and to give an insight into the behaviour of the star at the same time. Obviously, possibly superposed δ Scuti-type pulsations would be easily detectable as well.

3. Observation of γ Doradus stars with the main CAM

A star which is to be observed as a MONS prime target must be as bright as possible and its γ Dor nature must be well established. By examination of Table 1 we can see that 3 of the bona fide γ Dor stars have apparent magnitudes between 4 and 5, i.e. in the typical magnitude range of the solar-type variables included in MONS' prime program. Those are γ Dor, 9 Aur and BS 2740. The three stars have been observed at least through one successful multi-longitude multi-colour photometric campaign and the presence of NRP is established beyond any reasonable doubt. The cases of γ Dor and 9 Aur are particularly interesting since these are the most thoroughly observed and studied γ Dor stars so far. We now turn to short descriptions of these objects.

The bright star γ **Dor** was observed in the sixties during survey work dedicated to some chemically peculiar A stars and was suspected to show variations with an amplitude $\Delta V = 0^m.05$. Cousins (1992) observed it with high precision in 1983/84 (55 measurements in 120 nights). The analysis provided evidence of two close frequencies at 1.32 and $1.36\,\mathrm{d}^{-1}$, which was confirmed, together with their beat frequency at $0.04\,\mathrm{d}^{-1}$, by a further intensive observing run in 1989. The star has since then been object of two multi-longitude photometric and one multi-longitude spectroscopic campaign (Balona et al. 1994, 1996). The first campaign confirmed the two close frequencies at $f_1 = 1.32098$ and f_2=$1.36354\,\mathrm{d}^{-1}$ and provided evidence for the presence of a third frequency at f_3=$1.47447\,\mathrm{d}^{-1}$. The second campaign resulted in clear evidence of line profile variations due to NRP.

The availability of line profiles sampled with sufficient resolution allowed these authors to apply the *moment method* (Balona 1987, Aerts et al. 1992) to this star. Such a method constructs a *discriminant*, the minimum value of which indicates the most suitable combination of i, l, m to fit the observed oscillation. The discriminant diagrams for γ Dor computed by Balona et al. (1996) are reported in Figure 1 (right box). Clearly the results are non-conclusive since they do neither point to a unique solution nor, as they should, to a unique value for the inclination. However, Balona et al. (1996) state that the most probable angle of inclination for the star is $i = 70^o$ and the best estimate for the pulsational quantum numbers (l, m) are (3,3) for f_1 and (1,1) both for f_2 and f_3.

Photometric variability of the bright northern star **9 Aurigae** was reported by Krisciunas & Guinan (1990). The star was then observed by Krisciunas et al. (1995) through a multi-longitude campaign, however only comprising sites in the continental US and Hawaii. The analysis of these data resulted in the detection of two frequencies at $f_1 = 0.794$ and f_2=$0.345\,\mathrm{d}^{-1}$. Radial velocity measurements were also collected during the campaign with the CORAVEL instrument in France. Radial velocity curves gave evidence for variability with frequency f_2, but no sign of f_1, whereas



Fig. 1.— Left box: phased Hipparcos light curves of suspected γ Dor stars reported by Aerts, Eyer & Kestens (1998). Right box: results of the moment analysis of γ Dor presented by Balona et al. (1996). The figure shows the discriminants in arbitrary units as a function of the angle of inclination (in degrees) for various modes (l, m). f_1, f_2 and f_3 are represented in the top, mid and bottom panel, respectively.

the second output quantity of CORAVEL, the line width, was dominated by f_1, as the photometry. This prompted Aerts and Krisciunas (1996) to apply the moment method to the same CORAVEL data. These authors reported a possible identification of the two frequencies as the manifestation of an $l = 3$, $|m| = 1$ spheroidal mode and its toroidal correction due to stellar rotation.

In order to overcome the aliasing problem, 9 Aur was re-scheduled for a multi-longitude photometric campaign; the results were published by Zerbi et al. (1997). This time a coverage of more than 18^h over 24^h was achieved in a number of nights resulting in a substantial decrease of the $1\,\mathrm{d}^{-1}$ alias. The main frequency f_1 was confirmed and a new frequency at f_3=0.7681 d^{-1} was discovered well above the level of detection. However, the previously known mode at $f_2 = 0.3454$ could not be clearly resolved. Its power was distributed into two peaks at f_{2a}=0.2788d^{-1} and at $f_1/2$=0.3975 d^{-1}, the latter causing the phased curve of the main signal to be double-wave-shaped. Most of the measurements in this campaign were collected in the Strömgren system. This made it possible to observe how the oscillations manifested themselves in the c_1 index (tracing the luminosity) and the $(b - y)$ index (tracing the effective temperature) for the principal frequency. It was found that they occurred in phase within the errors. This means that geometrical displacements are not relevant for the oscillations, which can be assumed as due basically to temperature variations. To some extent this is what one would expect for a low-order g-mode oscillation.

The light variability of the F0 IV star **BS 2740** was discovered by Hensberge et al. (1981), who used it as a comparison star for Ap star observations. On the basis of 20 *uvby* measurements collected in 16 nights spanning 41 days, these authors derived a period of 0.936 d^{-1} and classified BS 2740 as a "mild Ap star". This evidence and the star's position in the HR diagram combined with the detection of line-profile variations in unpublished spectrograms prompted Poretti et al. (1997) to investigate its possible γ Dor nature through a multi-longitude campaign. The light curve of BS 2740 was explained with four frequencies, $f_1 = 1.0434$, $f_2 = 0.9951$, $f_3 = 1.1088$ and f_4=0.9019 d^{-1} and white noise. All the frequencies above are very close to 1 d^{-1}, which means that roughly the same phase of the oscillations would be seen by a single-site observer. Multisite campaigns or space observations are therefore required for this star.

4. Conclusions

The γ Dor stars are known to exhibit g-mode pulsations at time scales that make their observation difficult from the ground because of aliasing problems. The location of the class in the HR diagram and some theoretical arguments imply that solar-like oscillations could be excited in these stars as well. The MONS satellite is then the ideal tool to investigate both aspects taking profit of the clean spectral window of the STs and the precision of the CAM. We therefore strongly support the inclusion of a suitable candidate in the main target list and the recording of as many targets as possible with the STs.

REFERENCES

Aerts C., Eyer L., Kestens E., 1998, A&A, 337, 790

Aerts C., De Pauw M., Waelkens C., 1992, A&A, 266, 294

Aerts C., Krisciunas K., 1996, MNRAS, 278, p. 877

Balona L.A., 1987, MNRAS, 224, 41

Balona L.A., Krisciunas K., Cousins A.W.J. 1994, MNRAS, 270, 905

Balona L.A., Böhm T., Foing B. H., et al. 1996, MNRAS, 281, 1315

Cousins A.W.J. 1992, Observatory, 112, 53

Guzik, J.A., Kaye, A.B., Bradley, P.A., Cox, A.N., Neuforge, C., 2000, in "The Impact of Large-Scale surveys on Pulsationg Star Reasearch", IAU Coll. 176, Szabados and Kurtz eds., in preparation.

Handler G., 1999a, MNRAS, 309, L19

Handler G., 1999b, SAAO Newsletter 33, 1

Hensberge H., Deridder G., Doom C., et al. 1981, A&AS, 46, 151

Kaye A.B., Handler G., Krisciunas K., Poretti E., Zerbi F.M., 1999, PASP, 111, 840

Krisciunas K., Guinan E. F., 1990, IBVS 3511

Krisciunas K., Griffin R. F., Guinan E. F., Luedeke K. D., McCook G. P. 1995, MNRAS, 273, 662

Poretti E., Koen C., Martinez P., et al., 1997, MNRAS, 292, 621

Zerbi F.M., Garrido R., Rodríguez E., et al. 1997, MNRAS, 290, 401